

IgE Food Antibody Assessment



63 Zillicoa Street
Asheville, NC 28801
© Genova Diagnostics

Patient: **SAMPLE**
PATIENT

Age:

Sex:

MRN:

| IgE Food Antibody Results | | | |
|----------------------------------|----------------|-------|-----------|
| | RESULT kU/L | CLASS | INDICATOR |
| Grains | | | |
| Buckwheat | 0.89 | II | |
| Corn | 16.31 | V | |
| Oat | <0.24 | 0/1 | |
| Rice | <0.24 | 0/1 | |
| Sesame | <0.24 | 0/1 | |
| Soybean | <0.24 | 0/1 | |
| Wheat | 1.3 | III | |
| Dairy | | | |
| Egg White | 0.26 | I | |
| Cow's Milk | <0.24 | 0/1 | |
| Nuts | | | |
| Almond | 0.24 | 0/1 | |
| Brazil Nut | <0.24 | 0/1 | |
| Coconut | 0.4 | II | |
| Hazelnut | <0.24 | 0/1 | |
| Peanut | 98.36 | VI | |
| Seafood | | | |
| Blue Mussel | 26.12 | VI | |
| Codfish | 3.89 | III | |
| Salmon | <0.24 | 0/1 | |
| Shrimp | 3.9 | III | |
| Tuna | <0.24 | 0/1 | |

| Total IgE | | | |
|------------------|--------|--------------|-----------------|
| | Inside | Outside | Reference Range |
| Total IgE | | 139.1 | <=87.0 IU/mL |

- IgE levels must be used in conjunction with the clinical picture and are not intended to be independently diagnostic.
- The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. All assays are cleared by the U.S. Food and Drug Administration.
- Total IgE level may have clinical significance regardless of specific antibody levels.
- Increasing levels of antibody detected suggest an increasing clinical reactivity to specific foods.

| Key | | | |
|------------|-------------|---------------------------|-----------|
| Class | kU/L | Levels of Specific IgE | Indicator |
| 0/1 | <=0.24 | Undetectable or Equivocal | |
| I | 0.25 - 0.39 | Low | |
| II | 0.4 - 1.29 | Moderate | |
| III | 1.3 - 3.89 | High | |
| IV | 3.9 - 14.99 | Very High | |
| V | 15 - 24.99 | Very High | |
| VI | >=25 | Very High | |

| Laboratory Comments |
|----------------------------|
| |